



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Clinical outcomes among COVID-19 patients managed with modern and traditional Siddha medicine -A retrospective cohort study, Chennai, Tamil Nadu, India, 2020

Ramaswamy Meenakumari, Karuppiyah Thangaraj, Arunachalam Sundaram, Malayappan Meenakshi Sundaram, Ponnappan Shanmugapriya, Andi Mariappan, Melvin George, Venkatesan Suba, Elumalai Rajalakshmi, Muthappan Sendhilkumar

PII: S0975-9476(21)00118-2

DOI: <https://doi.org/10.1016/j.jaim.2021.06.010>

Reference: JAIM 470

To appear in: *Journal of Ayurveda and Integrative Medicine*

Received Date: 26 February 2021

Revised Date: 30 May 2021

Accepted Date: 14 June 2021

Please cite this article as: Meenakumari R, Thangaraj K, Sundaram A, Sundaram MM, Shanmugapriya P, Mariappan A, George M, Suba V, Rajalakshmi E, Sendhilkumar M, Clinical outcomes among COVID-19 patients managed with modern and traditional Siddha medicine -A retrospective cohort study, Chennai, Tamil Nadu, India, 2020, *Journal of Ayurveda and Integrative Medicine*, <https://doi.org/10.1016/j.jaim.2021.06.010>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 The Authors. Published by Elsevier B.V. on behalf of Institute of Transdisciplinary Health Sciences and Technology and World Ayurveda Foundation



Clinical outcomes among COVID-19 patients managed with modern and traditional Siddha medicine -A retrospective cohort study, Chennai, Tamil Nadu, India, 2020

*Ramaswamy Meenakumari^a, Karuppiiah Thangaraj^b, Arunachalam Sundaram^b, Malayappan Meenakshi Sundaram^a, Ponnappan Shanmugapriya^a, Andi Mariappan^a, Melvin George^b, Venkatesan Suba^a, Elumalai Rajalakshmi^c, Muthappan Sendhilkumar^c

^a National Institute of Siddha (NIS), Tambaram Sanatorium, Chennai, Tamil Nadu, India

^b SRM Medical College, Kattankulathur, Chennai, Tamil Nadu

^c ICMR-National Institute of Epidemiology (ICMR-NIE), Chennai, Tamil Nadu

Corresponding author

Dr Meenakumari Ramaswamy

National Institute of Siddha (NIS),
Tambaram Sanatoruim,
Chennai, Tamil Nadu 600047

Maid ID: mkumari474@yahoo.com

Clinical outcomes among COVID-19 patients managed with modern and traditional Siddha medicine -A retrospective cohort study, Chennai, Tamil Nadu, India, 2020

Abstract

Background

Kabasura Kudineer (KSK) is a Siddha polyherbal decoction recommended by the Ministry of AYUSH and the Tamil Nadu government to prevent and manage COVID-19 in India. In this context, we aimed to determine the outcome of integrated therapy for COVID-19 using KSK in virologic clearance and length of hospital stay.

Methods

It was a single-centre, retrospective cohort study. We included the COVID-19 patients admitted to SRM Medical College Hospital and Research Centre, Chennai, during May-June 2020. The KSK was administered along with the standard of care for all the patients. We collected demographic, clinical data and laboratory parameters data and were presented as frequencies and proportions.

Results

We collected 204 COVID-19 positive patients' data. The mean (SD) age was 39.5 (13.4) years with a range of 13-79. Majority of the patients were male (n=157; 77%), 28% (n=58) had any co-morbidities and 61% (n=131) were with mild symptoms. Fever (n=57; 27.9%) and cough (n=53; 25.9%) were the commonly reported symptoms. Paracetamol (n=135; 66.7%) and Zincovit (n=197, 96.6%) were the commonly administered medicines along with KSK. About 74% of asymptomatic (n=54) and 65% of mild symptomatic (n=85) patients turned negative for COVID-19 in RT-PCR within 4-7 days. There was a significant difference in the blood parameters ($p<0.05$) after the integrated treatment.

Conclusion

1 The use of KSK with standard care of treatment in COVID-19 treatment had notable results in
2 the duration taken for virologic clearance, thereby reducing the length of hospital stay and
3 improvement in laboratory parameters.

4 **Keywords:** Siddha, *Kabasura Kudineer*, COVID-19, AYUSH, Traditional medicine

1. Introduction

The Severe Acute Respiratory Syndrome-CoronaVirus-2 (SARS-CoV-2) infection became a global health threat to the community and severely affected livelihood. More than 100 million confirmed cases and 2 million deaths due to corona virus disease 2019 (COVID-19) are reported globally since it has been identified in December 2019 [1]. However, most people with COVID-19 experience mild and moderate symptoms and approximately 10-15% of cases progress to severe illness [2,3]. Currently, there are no treatments due to the novelty of the virus and its broad clinical spectrum. In the absence of effective therapies, traditional medicines were used as an integrated treatment for managing COVID-19, as happened in the past with other infectious diseases [4].

In India, Ayurveda, Yoga and Naturopathy, Unani, Siddha and Sowa-Rigpa and Homoeopathy (AYUSH) are the traditional medicines. Siddha system of medicine is primarily practiced in Southern parts of India and Tamil speaking regions of the world. In response to the COVID-19 pandemic, the Ministry of AYUSH of India has issued treatment guidelines for AYUSH practitioners to prevent and manage COVID-19, including the administration of *Kabasura Kudineer* (KSK), a polyherbal Siddha formulation [5]. The health ministry approved the decision based on the experience of administering polyherbal Siddha formulations in the management of Chikungunya and Dengue outbreak in the state of Tamil Nadu [6,7].

The clinical features of SARS-CoV-2 infection may be compared with *Kabasuram* (*Iya suram*) as per the Siddha literature, which is basically due to elevation of *Iyam*, one of the three humors of the body. Elevated *Iyam* and Vali and Azhal become Thontham (all three humors mixed and elevated), which leads to *sanni* which is defined as a severe and fatal stage of a disease [8]. KSK a polyherbal Siddha formulation which has been generally prescribed and administered to manage *Iyasuram*, characterized with fever and flu like symptoms by Siddha medical practitioners [9]. KSK comprises 15 herbs [10]. The Ministry of AYUSH has recommended in the advisory guidelines for the prophylaxis and management of COVID-19 based on the clinical experience and evidence of pharmacological, phytochemical activities of its ingredients. However, it showed antipyretic, anti-viral and immunomodulatory properties in initial phytochemical, pharmacological and *in-vitro* studies [11].

The National Institute of Siddha (NIS) which is located in Chennai, Tamil Nadu with its excellence in research and higher education in Siddha system of medicine and played a pivotal role in COVID-19 response. NIS has involved in COVID-19 response activities in collaboration with medical, academic and research institutions in Tamil Nadu [12]. The SRM Medical college Hospital and Research Centre, a tertiary care centre in Tamil Nadu, is one such health facility in which the KSK was administered along with standard care of treatment for COVID-19 patients under the supervision of Siddha faculty of NIS as a collaborative activity. This polyherbal decoction is used in general practice in Siddha system. Though it is yet to be demonstrated by controlled clinical trials for its safety and efficacy, we got the opportunity to document the experience of administering KSK along with the routine care at SRM health facility. During the pandemic, KSK has been advocated to administer as a preventive measure and the standard care for asymptomatic, mild and moderate symptomatic COVID-19 patients based on the evidence showing anti-viral properties in the preliminary pharmacological and phytochemical studies [11]. However, clinical documentation is highly needed to know the benefit of integrated treatment. In this context, we aimed to determine the clinical outcome of integrated therapy for COVID-19 using *Kabasura kudineer* in terms of virologic clearance for COVID-19 in RT-PCR and to estimate the length of hospital stay.

2. Methods

2.1. Study setting

The study site is SRM Medical College Hospital and Research Centre, Chennai, Tamil Nadu. This is a tertiary care hospital which serves around 3, 00,000 populations in the surrounding areas.

2.2. Study participants

COVID-19 positive patients who were admitted in SRM Medical College Hospital and treated with integrated therapy.

2.3. Inclusion and exclusion criteria

All patients included in this study were diagnosed for COVID-19 based on the World Health Organization (WHO) guidelines those who had influenza like illness and were SARS-CoV-2 Reverse transcription polymerase chain reaction (RT-PCR) positive in throat swab. We included

consecutive COVID-19 positive patients with asymptomatic and mild symptoms irrespective of age and gender who were admitted during the study period and treated with KSK along with the standard care of treatment.

2.4. Study design

This was a single-centre retrospective cohort study.

2.5. Study period:

The COVID-19 patients admitted during May – June 2020 were included in this study.

2.6. Operational definitions

2.6.1. Virologic clearance

The time duration is taken from the first SARS-CoV-2 virus nucleic acid test result positive to the first negative result. Throat swabs of patients were taken to test if the patients were relieved from all the clinical symptoms for consecutive 3 days with normal oxygen saturation.

2.6.2. Length of Hospital stay

The time duration is taken from hospital admission to the discharge for a patient.

2.7. Standard care of treatment

All the COVID-19 patients were managed symptomatically with the administration of antipyretics, antibiotics, expectorant, Zinc and Vitamin C and the regular treatment related to their comorbid conditions such as anti-diabetic and anti-hypertensive drugs.

2.8. Siddha system management with *Kabasura Kudineer* (KSK)

The KSK is a polyherbal formulation comprising of 15 medicinal herbs (**Table 1**). The medicine was purchased from a Good Manufacturing Practice (GMP) certified company in order to ensure the good quality. The decoction was prepared by adding 10 gms of KSK powder with 240 ml of water, boiled and reduced to one-fourth (60 ml) and filtered. The decoction (60 ml) was administered orally thrice daily, 20 min before food for all the patients. Each dose of KSK decoction was administered to all the patients within 3 hours of preparation. The pharmacist of

NIS has prepared the decoction to ensure the quality of medicine. The clinical and paramedical SRM hospital staff ascertained that the medicine was administered to the patients as per the treatment protocol prescribed. First dose of KSK was administered within 24 hrs of admission and continued thrice daily throughout the hospital stay. The patients were stayed in the hospital until they get the virologic clearance by RT-PCR test. Further, they were advised to continue KSK twice daily for at-least a week after discharge.

2.9. Data collection methods

We used data abstraction forms to retrieve information from the medical records. The information collected are the demographic details such as age gender, hospitalization details such as date of admission and discharge, medical history for comorbid conditions, vital parameters, all clinical signs and symptoms including other than for COVID-19. The laboratory parameters of blood test, RT-PCR findings (initial date of positive and the date when the patient became negative for COVID-19 in RT-PCR) and the medical interventions underwent such as KSK administration and the list of drugs administered under standard treatment of care also collected. We included all the patients admitted and treated in the study period, irrespective of the date of laboratory confirmation and the date of admission to the hospital. Hence, there may be a time lag between the date of first RT PCR test on which they found positive and the date of admission in the hospital for some patients. The decoction of KSK was administered three times in a day to all the asymptomatic and patients with mild symptoms. The KSK was administered irrespective of the presence of any co-morbid condition. Frequency and dose of administration of KSK were also abstracted from the medical records. Throat swabs of patients were taken for testing if the patients were relieved from all the clinical symptoms for continuous 3 days with normal oxygen saturation

2.10. Statistical analysis

The collected data were entered in excel and cleaned before analysis. Continuous variables were described as means and SD, or as median and range and the categorical variables were expressed as frequency and proportions. Paired t-test was done to find out the significant difference in any paired variables. All statistical analyses were performed using SPSS version 16.0.

2.11. Ethics approval

This study was approved by the Institutional Ethics Committee of SRM Medical College hospital and research centre, Chennai (EC No. 1968/IEC/2020) to access and analyse the hospital records. Since we reviewed only the hospital records, obtaining informed consent from the participants was waived by the Institutional Ethics Committee. We did not use any personal identifiers during the analysis and not revealed anywhere in this report.

3. Results

3.1. Baseline profile of the study participants

We collected 204 confirmed COVID-19 patients' data. The mean (SD) age was 39.5 (13.4) years with a range of 13-79. Majority of the patients were male (n=157; 77%), and 28% (n=58) had any of the co-morbid conditions. Among the comorbidities, diabetes (n=35; 17.2%) and hypertension (n=26; 12.7%) were the predominantly reported. The mean value of systolic and diastolic blood pressure was 119.1 (11.2%) mm Hg and 77.9 (7.9%) mm Hg, respectively on the day of admission. The median hospital stay was 9 days (IQR 10-7) (**Table 2**).

3.2. COVID-19 related status of the included study participants

Among the total, 131(64%) were with mild symptoms of COVID-19. The most common clinical symptoms reported by the patients were fever 57 (27.9%), cough 53 (25.9%), sore throat 29 (14.2%), headache 24 (11.7%) and fatigue 13(6.4%) during the hospital stay (**Figure 1**).

3.3. Clinical outcomes among the study participants

Most of the asymptomatic (n=54; 74%) patients turned negative for COVID-19 in RT-PCR within 4-7 days and 25% (n=18) in 8-15 days after integrated treatment with *Kabasura kudineer*. Whereas for mild symptomatic it was 65% (n=85) and 34% (n=44) in 4-7 days and 8-15 days respectively. Nearly two-thirds of Hypertension patients (n=17; 65.4%) turned negative in 4-7 days, whereas only half of the diabetes patients (n=20; 57.1%) turned negative in 4-7 days (**Table 3**). Kaplan-Meier curve (**Figure 2**) shows the median number of days taken for RT-PCR negative from the date of admission based on the presence and absence of comorbid condition is 7 days (CI 6.783-7.217). The log rank test showed the two curves were not significantly different (P=0.895). However, there was a significant difference among patients in the counts of total

lymphocytes, eosinophils and basophils ($p<0.05$) (**Table 4**) counts before and after integrated treatment. Among the total, there was one patient died who had presented with comorbid conditions.

3.4. Adverse drug reactions

No significant adverse reactions were recorded except diarrhoea ($n=4$; 2%) and gastritis ($n=3$; 1.5%) during the hospital stay (**Table 2**).

3.5. Drug profile

Antipyretic such as Paracetamol ($n=136$; 66.7%), antibiotics such as Azithromycin ($n=76$; 37.3%), expectorants such as syrup Ascoril ($n=36$; 17.6%) and supplementary of Zincovit ($n=197$; 96.6%) were the conventional drugs used during the integrated treatment (**Table 5**). None was withdrawn from the integrated treatment due to treatment-related discomfort or any adverse reactions.

4. Discussion

The WHO declared the ongoing outbreak as a global public health emergency [13]. There are no specific antiviral drugs that are effective against COVID-19 at present [14]. COVID-19 caused by SARS-CoV-2 has spread rapidly, with the characteristics of high infectivity and high susceptibility [15]. In the early phases of COVID-19 pandemic, management using Siddha herbal formulations and the routine care of treatment was tried in the South Indian setting. The NIS, an academic and research institution had an opportunity to collaborate with the existing state government health facilities and other academic and research institutions. It paved the way to administer selected polyherbal formulations as per the proposed treatment guidelines by the Ministry of AYUSH and observe the outcomes. Hence, we were able to document the clinical experience in the management of COVID-19 using KSK combined with the routine care of treatment.

COVID-19 patients often carry SARS-CoV-2 for a longer time; hence prolonged antiviral treatment is needed to become negative in virology test. This duration is a concern in the prognosis of disease, especially among the elderly and patients with severe symptoms. Patients with chronic SARS-CoV-2 infection may develop severe lung infections and acute respiratory distress syndrome, leading to hospitalization. The COVID-19 pandemic has put unprecedented

pressure on the health care system. Hence, it is crucial to shorten the duration of hospital stay of the patients. As per the revised discharge protocol issued by the Ministry of Health and Family Welfare (MoHFW), the Government of India Revised Discharge Policy for COVID-19, May 2020, the length of hospital stay in standard allopathy care of mild cases was 10 days ([16]. In the current observation, around two-thirds of the mild symptomatic patients were relieved from symptoms, became negative for RT-PCR test, and discharged from the hospital in 10 days, which is as mentioned in the discharge policy issued by the MoHFW. The course of integrated treatment with KSK ranged from 4-18 days irrespective of the symptom status, which is shorter than the studies reported earlier from China (4-53 days) and from other countries (4-21 days) outside China ([17].

The integrated treatment shortens the time taken to become negative for COVID-19 and thereby reducing the length of hospital stay. With these observations, it may be postulated that if KSK is administered with the routine treatment, it will benefit the effective management of COVID-19. People with known diabetes and hypertension and older age group are at a greater risk of contracting infection, worse prognosis and mortality with respect to COVID-19 infection [18,19]. In contrary the time taken for virologic clearance was 5-15 days for both the group who had either diabetes or hypertension in the current observation.

Lymphopenia is reported in severe cases and the patients with poor prognosis [20,21]. During the current integrated therapy with KSK the identified patients with low count of lymphocytes, eosinophils and basophils at the time of admission have shown significant improvement at discharge. It is notable to mention that many of the phytoconstituents of KSK formulation are studied for their immunomodulatory potential and activities [22,23]. The ingredients in the selected Siddha formulation KSK has anti-pyretic, anti-inflammatory and immunomodulatory activities, which can appropriately suitable for the management of COVID-19 [11, 22-30]. Research studies are conducted to analyse phytochemicals from KSK against the SARS-CoV-2 through structure-based *in silico* molecular docking and identified potent anti-COVID-19 natural compounds [31,32]. Furthermore, clinical trials are also implemented to determine the efficacy of KSK in comparison with other standard treatment protocol [33].

Limitations

This study has few limitations: (i) No randomization involved, thereby leading to selection bias. However, we have included all the patients admitted during the study period. (ii) There was no comparison group in this study due to ethical consideration in treating the novel infection in the absence of standard treatment or cure. It resulted in limiting the scope of establishing analytical evidence of the integrated treatment.

Conclusion

In conclusion, the integrated management of COVID-19 with KSK and standard care of treatment has shown notable results in virologic clearance, thereby reducing hospital stay length than the discharge policy issued by health authorities. Moreover, there were no significant adverse reactions concerning the administration of KSK.

Recommendations

We recommend conducting studies with a comparator group to determine the significant difference in the outcome and randomized controlled trials to evaluate the safety and efficacy of KSK.

Declaration of competing interests

The authors declare that there is no competing interest.

Acknowledgements

Our sincere thanks to the ministry of AYUSH for supporting clinical research projects of Siddha system of medicine for COVID-19. We thank the frontline medical staff of SRM Medical College and Research Institute for their involvement and contribution in the study.

References:

1. World Health Organization. WHO Coronavirus Disease (COVID-19) Dashboard. [Accessed on 20 February 2021]. Available from: <https://covid19.who.int/>
2. World Health Organization. What we know about Long-term effects of COVID-19. [Accessed on 20 February 2021]. Available from: https://www.who.int/docs/default-source/coronaviruse/risk-comms-updates/update-36-long-term-symptoms.pdf?sfvrsn=5d3789a6_2
3. Alschuler L, Chiasson AM, Horwitz R, Sternberg E, Crocker R, Weil A, Maizes V. Integrative medicine considerations for convalescence from mild-to-moderate COVID-19 disease. Explore. 2020 Dec 23.
4. Lee MS, Song E. Integrative medicine for COVID-19: researches and evidence. Integrative Medicine Research. 2020 Aug 4. 2020 Sep; 9(3)
5. Ministry of AYUSH. [Accessed on Jan 25, 2021]. Available from: <https://www.ayush.gov.in/ayush-guidelines.html>
6. Government of Tamil Nadu. Directorate of Public Health & Preventive Medicine. [Accessed Jan 25, 2021]. Available from: <https://www.tnhealth.org/dphfacts/chikungunya.htm>.
7. Government of Tamil Nadu . Health & Family Welfare Department. [Accessed on Jan 25, 2021]. Available from: <https://tnhealth.tn.gov.in/tngovin/dph/dphdbdengue.php>.
8. Thiagarajan R. Yugimunivar vaidhyachinthamani (Perunool 800, Tamil). 2nd ed. Chennai, Tamil Nadu: Palani Thandayuthapani Devasthanam publications, Directorate of Indian systems of Medicine; 1976.

9. Kuppusamy Mudaliar KN. Siddha Maruthuvam (Tamil). Chennai: The Directorate of Indian Medicine and Homoeopathy, Govt. Of Tamil Nadu; 1987.
10. Kuppusamy Mudaliar KN, Uthamarayan KS. Siddha Vaidhya Thirattu Chennai: The Directorate of Indian Medicine and Homoeopathy, Govt. Of Tamil Nadu; 2009. p.293-294.
11. Government of Tamil Nadu. The Tamil Nadu Dr. M.G.R. Medical University. . [Accessed on Feb 6, 2021]. Available from: <http://repository-tnmgrmu.ac.in/5357/>
12. National Institute of Siddha. [Accessed on Feb 6, 2021]. Available from: <https://nischennai.org/index.html>
13. World Health Organization. Clinical management of severe acute respiratory infection when novel coronavirus (2019-nCoV) infection is suspected: interim guidance, 28 January 2020. Available from. <https://apps.who.int/iris/handle/10665/330893>
14. World Health Organization. Statement on the second meeting of the International Health Regulations (2005) Emergency Committee regarding the outbreak of novel coronavirus (2019-nCoV). Available from: [https://www.who.int/news/item/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-\(2005\)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-\(2019-ncov\)](https://www.who.int/news/item/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-(2005)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-(2019-ncov))
15. Sohrabi C, Alsafi Z, O'Neill N, Khan M, Kerwan A, Al-Jabir A, Iosifidis C, Agha R. World Health Organization declares global emergency: A review of the 2019 novel corona virus (COVID-19). International journal of surgery. 2020 Apr 1;76:71-6.
16. Ministry of Health and Family welfare. Revised Discharge Policy for COVID-19. [Accessed on Feb 9 2021]. Available from: <https://www.mohfw.gov.in/pdf/ReviseddischargePolicyforCOVID19.pdf>
17. Rees EM, Nightingale ES, Jafari Y, Waterlow NR, Clifford S, Pearson CA, Jombart T, Procter SR, Knight GM, CMMID Working Group. COVID-19 length of hospital stay: a systematic review and data synthesis. BMC medicine. 2020 Dec;18(1):1-22.
18. Rodríguez-Molinero A, Gálvez-Barrón C, Miñarro A, Macho O, López GF, Robles MT, Dapena MD, Martínez S, MilàRàfols N, Monaco EE, Hidalgo García A. Association between COVID-19 prognosis and disease presentation, comorbidities and chronic treatment of hospitalized patients. PloS one. 2020 Oct 15;15(10):e0239571.
19. Kluge H. Older people are at highest risk from COVID-19, but all must act to prevent

- community spread. World Health Organization for Europe. 2020.
20. Fathi N, Rezaei N. Lymphopenia in COVID-19: Therapeutic opportunities. *Cell biology international*. 2020 Sep;44(9):1792-7.
 21. Huang I, Pranata R. Lymphopenia in severe coronavirus disease-2019 (COVID-19): systematic review and meta-analysis. *Journal of intensive care*. 2020 Dec;8:1-0.
 22. Dwivedi SK, Enespa A. *Tinosporacordifolia* with reference to biological and microbial properties. *International Journal of Current Microbiology and Applied Sciences*. 2016;5(6):446-65.
 23. Hossain MD, Urbi Z, Sule A, Rahman KM. *Andrographis paniculata* (Burm. f.) Wall. ex Nees: a review of ethnobotany, phytochemistry, and pharmacology. *The Scientific World Journal*. 2014 Jan 1;2014.
 24. Soltan MM, Zaki AK. Antiviral screening of forty-two Egyptian medicinal plants. *Journal of ethnopharmacology*. 2009 Oct 29;126(1):102-7.
 25. Pawar VA, Pawar PR. *Costus speciosus*: an important medicinal plant. *International journal of science and Research*. 2014 Jul;3(7):28-33.
 26. Gurgel AP, da Silva JG, Grangeiro AR, Oliveira DC, Lima CM, da Silva AC, Oliveira RA, Souza IA. In vivo study of the anti-inflammatory and antitumor activities of leaves from *Plectranthus amboinicus* (Lour.) Spreng (Lamiaceae). *Journal of Ethnopharmacology*. 2009 Sep 7;125(2):361-3.
 27. Batiha GE, Alkazmi LM, Wasef LG, Beshbishy AM, Nadwa EH, Rashwan EK. *Syzygium aromaticum* L.(Myrtaceae): Traditional uses, bioactive chemical constituents, pharmacological and toxicological activities. *Biomolecules*. 2020 Feb;10(2).
 28. Manouze H, Bouchatta O, Gadhi AC, Bennis M, Sokar Z, Ba-M'hamed S. Anti-inflammatory, antinociceptive, and antioxidant activities of methanol and aqueous extracts of *Anacyclus pyrethrum* roots. *Frontiers in pharmacology*. 2017 Sep 5;8:598.
 29. Kesharwani A, Polachira SK, Nair R, Agarwal A, Mishra NN, Gupta SK. Anti-HSV-2 activity of *Terminalia chebula* Retz extract and its constituents, chebulagic and chebulinic acids. *BMC Complement Altern Med*. 2017 Feb 14;17(1):110.
 30. Sood R, Raut R, Tyagi P, Pareek PK, Barman TK, Singhal S, Shirumalla RK, Kanoje V, Subbarayan R, Rajerethinam R, Sharma N. *Cissampelos pareira* Linn: Natural source of potent antiviral activity against all four dengue virus serotypes. *PLoS neglected tropical*

diseases. 2015 Dec 28;9(12):

31. Vincent S, Arokiyaraj S, Saravanan M, Dhanraj M. Molecular Docking Studies on the Anti-viral Effects of Compounds From Kabasura Kudineer on SARS-CoV-2 3CLpro. *Frontiers in molecular biosciences*. 2020;7.
32. Kiran G, Karthik L, Devi MS, Sathiyarajeswaran P, Kanakavalli K, Kumar KM, Kumar DR. In silico computational screening of Kabasura Kudineer-official Siddha formulation and JACOM against SARS-CoV-2 spike protein. *Journal of Ayurveda and integrative medicine*. 2020 May 25.
33. Natarajan S, Anbarasi C, Sathiyarajeswaran P, Manickam P, Geetha S, Kathiravan R, Prathiba P, Pitchiahkumar M, Parthiban P, Kanakavalli K, Balaji P. The efficacy of Siddha Medicine, Kabasura Kudineer (KSK) compared to Vitamin C & Zinc (CZ) supplementation in the management of asymptomatic COVID-19 cases: A structured summary of a study protocol for a randomised controlled trial. *Trials*. 2020 Dec;21(1):1-2.

1

Journal Pre-proof

References:

1. World Health Organization. WHO Coronavirus Disease (COVID-19) Dashboard. [Accessed on 20 February 2021]. Available from: <https://covid19.who.int/>
2. World Health Organization. What we know about Long-term effects of COVID-19. [Accessed on 20 February 2021]. Available from: https://www.who.int/docs/default-source/coronaviruse/risk-comms-updates/update-36-long-term-symptoms.pdf?sfvrsn=5d3789a6_2
3. Alschuler L, Chiasson AM, Horwitz R, Sternberg E, Crocker R, Weil A, Maizes V. Integrative medicine considerations for convalescence from mild-to-moderate COVID-19 disease. *Explore*. 2020 Dec 23.
4. Lee MS, Song E. Integrative medicine for COVID-19: researches and evidence. *Integrative Medicine Research*. 2020 Aug 4. 2020 Sep; 9(3)
5. Ministry of AYUSH. [Accessed on Jan 25, 2021]. Available from: <https://www.ayush.gov.in/ayush-guidelines.html>
6. Government of Tamil Nadu. Directorate of Public Health & Preventive Medicine. [Accessed Jan 25, 2021]. Available from: <https://www.tnhealth.org/dphfacts/chikungunya.htm>.
7. Government of Tamil Nadu . Health & Family Welfare Department. [Accessed on Jan 25, 2021]. Available from: <https://tnhealth.tn.gov.in/tngovin/dph/dphdbdengue.php>.
8. Thiyagarajan R. Yugimunivar vaidhyachinthamani (Perunool 800, Tamil). 2nd ed. Chennai, Tamil Nadu: Palani Thandayuthapani Devasthanam publications, Directorate of Indian systems of Medicine; 1976.
9. Kuppusamy Mudaliar KN. Siddha Maruthuvam (Tamil). Chennai: The Directorate of Indian Medicine and Homoeopathy, Govt. Of Tamil Nadu; 1987.
10. Kuppusamy Mudaliar KN, Uthamarayan KS. Siddha Vaidhya Thirattu Chennai: The Directorate of Indian Medicine and Homoeopathy, Govt. Of Tamil Nadu; 2009. p.293-294.
11. Government of Tamil Nadu. The Tamil Nadu Dr. M.G.R. Medical University. . [Accessed on Feb 6, 2021]. Available from: <http://repository-tnmgrmu.ac.in/5357/>
12. National Institute of Siddha. [Accessed on Feb 6, 2021]. Available from: <https://nischennai.org/index.html>

13. World Health Organization. Clinical management of severe acute respiratory infection when novel coronavirus (2019-nCoV) infection is suspected: interim guidance, 28 January 2020. Available from: <https://apps.who.int/iris/handle/10665/330893>
14. World Health Organization. Statement on the second meeting of the International Health Regulations (2005) Emergency Committee regarding the outbreak of novel coronavirus (2019-nCoV). Available from: [https://www.who.int/news/item/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-\(2005\)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-\(2019-ncov\)](https://www.who.int/news/item/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-(2005)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-(2019-ncov))
15. Sohrabi C, Alsafi Z, O'Neill N, Khan M, Kerwan A, Al-Jabir A, Iosifidis C, Agha R. World Health Organization declares global emergency: A review of the 2019 novel corona virus (COVID-19). *International journal of surgery*. 2020 Apr 1;76:71-6.
16. Ministry of Health and Family welfare. Revised Discharge Policy for COVID-19. [Accessed on Feb 9 2021]. Available from: <https://www.mohfw.gov.in/pdf/ReviseddischargePolicyforCOVID19.pdf>
17. Rees EM, Nightingale ES, Jafari Y, Waterlow NR, Clifford S, Pearson CA, Jombart T, Procter SR, Knight GM, CMMID Working Group. COVID-19 length of hospital stay: a systematic review and data synthesis. *BMC medicine*. 2020 Dec;18(1):1-22.
18. Rodríguez-Molinero A, Gálvez-Barrón C, Miñarro A, Macho O, López GF, Robles MT, Dapena MD, Martínez S, MilàRàfols N, Monaco EE, Hidalgo García A. Association between COVID-19 prognosis and disease presentation, comorbidities and chronic treatment of hospitalized patients. *PloS one*. 2020 Oct 15;15(10):e0239571.
19. Kluge H. Older people are at highest risk from COVID-19, but all must act to prevent community spread. World Health Organization for Europe. 2020.
20. Fathi N, Rezaei N. Lymphopenia in COVID-19: Therapeutic opportunities. *Cell biology international*. 2020 Sep;44(9):1792-7.
21. Huang I, Pranata R. Lymphopenia in severe coronavirus disease-2019 (COVID-19): systematic review and meta-analysis. *Journal of intensive care*. 2020 Dec;8:1-0.
22. Dwivedi SK, Enespa A. *Tinosporacordifolia* with reference to biological and microbial properties. *International Journal of Current Microbiology and Applied Sciences*. 2016;5(6):446-65.
23. Hossain MD, Urbi Z, Sule A, Rahman KM. *Andrographis paniculata* (Burm. f.) Wall. ex

- Nees: a review of ethnobotany, phytochemistry, and pharmacology. The Scientific World Journal. 2014 Jan 1;2014.
24. Soltan MM, Zaki AK. Antiviral screening of forty-two Egyptian medicinal plants. Journal of ethnopharmacology. 2009 Oct 29;126(1):102-7.
 25. Pawar VA, Pawar PR. *Costus speciosus*: an important medicinal plant. International journal of science and Research. 2014 Jul;3(7):28-33.
 26. Gurgel AP, da Silva JG, Grangeiro AR, Oliveira DC, Lima CM, da Silva AC, Oliveira RA, Souza IA. In vivo study of the anti-inflammatory and antitumor activities of leaves from *Plectranthus amboinicus* (Lour.) Spreng (Lamiaceae). Journal of Ethnopharmacology. 2009 Sep 7;125(2):361-3.
 27. Batiha GE, Alkazmi LM, Wasef LG, Beshbishy AM, Nadwa EH, Rashwan EK. *Syzygium aromaticum* L.(Myrtaceae): Traditional uses, bioactive chemical constituents, pharmacological and toxicological activities. Biomolecules. 2020 Feb;10(2).
 28. Manouze H, Bouchatta O, Gadhi AC, Bennis M, Sokar Z, Ba-M'hamed S. Anti-inflammatory, antinociceptive, and antioxidant activities of methanol and aqueous extracts of *Anacyclus pyrethrum* roots. Frontiers in pharmacology. 2017 Sep 5;8:598.
 29. Kesharwani A, Polachira SK, Nair R, Agarwal A, Mishra NN, Gupta SK. Anti-HSV-2 activity of *Terminalia chebula* Retz extract and its constituents, chebulagic and chebulinic acids. BMC Complement Altern Med. 2017 Feb 14;17(1):110.
 30. Sood R, Raut R, Tyagi P, Pareek PK, Barman TK, Singhal S, Shirumalla RK, Kanoje V, Subbarayan R, Rajerethinam R, Sharma N. *Cissampelos pareira* Linn: Natural source of potent antiviral activity against all four dengue virus serotypes. PLoS neglected tropical diseases. 2015 Dec 28;9(12):
 31. Vincent S, Arokiyaraj S, Saravanan M, Dhanraj M. Molecular Docking Studies on the Anti-viral Effects of Compounds From Kabasura Kudineer on SARS-CoV-2 3CLpro. Frontiers in molecular biosciences. 2020;7.
 32. Kiran G, Karthik L, Devi MS, Sathiyarajeswaran P, Kanakavalli K, Kumar KM, Kumar DR. In silico computational screening of Kabasura Kudineer-official Siddha formulation and JACOM against SARS-CoV-2 spike protein. Journal of Ayurveda and integrative medicine. 2020 May 25.
 33. Natarajan S, Anbarasi C, Sathiyarajeswaran P, Manickam P, Geetha S, Kathiravan R,

Prathiba P, Pitchiahkumar M, Parthiban P, Kanakavalli K, Balaji P. The efficacy of Siddha Medicine, Kabasura Kudineer (KSK) compared to Vitamin C & Zinc (CZ) supplementation in the management of asymptomatic COVID-19 cases: A structured summary of a study protocol for a randomised controlled trial. *Trials*. 2020 Dec;21(1):1-2.

Table 1: Composition and ingredients of *Kabasura kudineer* administered with the standard care of treatment for COVID-19 during May-June 2020 at SRM Medical College Hospital and Research Centre

Name of herb in Tamil	Botanical Name	Quantity*
<i>Chukku</i>	<i>Zingiber officinale</i>	35 g
<i>Thippili</i>	<i>Piper longum</i>	35 g
<i>Lavangam</i>	<i>Syzygium aromaticum</i>	35 g
<i>Sirukaanjuriver</i>	<i>Tragia involucrata</i>	35 g
<i>Aragaram</i>	<i>Anacyclus pyrethrum</i>	35 g
<i>Karimulli</i>	<i>Solanum erianthum</i>	35 g
<i>Kadukkaithol</i>	<i>Terminalia chebula</i>	35 g
<i>Adathodaiilai</i>	<i>Justicia adhatoda</i>	35 g
<i>Karpooravalliilai</i>	<i>Coleus amboinicus</i>	35 g
<i>Koshtam</i>	<i>Sassure alappa</i>	35 g
<i>Seenthil</i>	<i>Tinospora cordifolia</i>	35 g
<i>Nilavembu</i>	<i>Andrographis paniculata</i>	35 g
<i>Vattathiruppi</i>	<i>Cissampelos pareira</i>	35 g
<i>.Koraikizhangu</i>	<i>Cyperus rotundus</i>	35 g
<i>Siruthekku</i>	<i>Clerodendrum serratum</i>	35 g

*Each dose is composed of a mixture of fifteen medicinal herbs in equal quantity.

Table 2. Demographic and clinical characteristics of patients with COVID-19 treated during May – June 2020 (n=204) in SRM Medical College Hospital and Research centre

Characteristics	Values	%
Age (years), mean \pm SD	39.53 \pm 13.4	
Gender		
Male	157	76.9
Female	47	23.0
Blood pressure		
Systolic (mean \pm SD)	119.1 \pm 11.2	
Diastolic (mean \pm SD)	77.9 \pm 7.9	
Hospital stay, [median (IQR)]	9 (7-10)	
Symptoms status		
Asymptomatic	73	36
Mild Symptomatic	131	64
Presence of Comorbidity	58	28.4
Co-morbid conditions		
Diabetes	35	17.2
Hypertension	26	12.7
Hypothyroidism	5	2.4
Renal Disease	2	1.0
Tuberculosis	1	0.5
Acid peptic disease	1	0.5
Septic arthritis	1	0.5
Microcytic anaemia	1	0.5
Dyslipidaemia	1	0.5
Coronary artery disease	1	0.5
Bronchial Asthma	1	0.5
Adverse drug events		
Diarrhoea	4	2.0
Gastritis/Gastroenteritis	3	1.5

Table 3. Frequency of patients turned negative for COVID-19 and the duration of integrated treatment using KSK during May – June 2020 (n=204) in SRM Medical College Hospital and Research centre

Symptomatic status	Duration of integrated treatment	Percentage of patients turned negative for RT-PCR
Asymptomatic	4-7 days	54 (74.0%)
	8-15 days	18 (24.7%)
	>15 days	1 (1.9%)
		N=73 (100%)
Mild symptomatic	4-7 days	85 (64.9%)
	8-15 days	44 (33.6%)
	>15 days	2 (1.5%)
		N=131 (100%)
Co-morbidities		
Diabetes Mellitus (Type 2)	4-7 days	20 (57.1%)
	8-15 days	16 (42.9%)
Hypertension	4-7 days	17 (65.4%)
	8-15 days	9 (34.6%)

Table 4: Difference in the blood parameters before and after the integrated treatment during May – June 2020 (n=204) in SRM Medical College Hospital and Research centre


Blood parameters	T	Df	Sig. (2-tailed)
Neutrophil	-1.356	13	.198
Lymphocyte	-5.479	10	.000
Eosinophil	-4.870	21	.000
Basophil	-3.138	49	.003
Monocyte	-.692	1	.614

1 **Table 5: Frequency of drugs used in the standard care to treat COVID-19 during May –**
 2 **June 2020 (n=204) in SRM Medical College Hospital and Research centre**

Nature of the drug	Medicine name	Number	%
Antipyretic	Paracetamol	136	66.7
Antibiotics	Azithromycin	76	37.3
	Cefoperazone	1	0.5
	Oflaxacin	1	0.5
	Inj. Cefoperazone sulbactam	1	0.5
	Vancomycin	1	0.5
	Ciproflaxacin	1	0.5
Supplementary	Zincovit	197	96.6
	Limcee	132	64.7
	Vit-c	85	41.7
	Becosules	56	27.5
	Bifilac	11	5.4
	Calcium	1	0.5
	Iron	1	0.5
Expectorants and antihistamines	Ascoril syrup	36	17.6
	Lecope	5	2.5
	Cetirizine/levocetirizine	7	3.4


3

Figure 1. Distribution of clinical signs and symptoms among COVID-19 patients admitted during May – June 2020 (n=204) in SRM Medical College Hospital and Research Centre, Chennai, Tamil Nadu



*One patient may have more than one clinical symptom

Figure 2: Kaplan-Meier for time taken for RT-PCR test negative based on the presence of any comorbid condition during May – June 2020 in SRM Medical College Hospital and Research Centre, Chennai, Tamil Nadu



Declaration of Conflict of Interest

Authors declare No financial or commercial conflict of interest.

Clinical outcomes among COVID-19 patients managed with modern and traditional Siddha medicine -A retrospective cohort study, Chennai, Tamil Nadu, India, 2020

*Ramaswamy Meenakumari^a, Karuppiath Thangaraj^b, Arunachalam Sundaram^b, Malayappan Meenakshi Sundaram^a, Ponnappan Shanmugapriya^a, Andi Mariappan^a, Melvin George^b, Venkatesan Suba^a, Elumalai Rajalakshmi^c, Muthappan Sendhilkumar^c

^a National Institute of Siddha (NIS), Tambaram Sanatorium, Chennai, Tamil Nadu, India

^b SRM Medical College, Kattankulathur, Chennai, Tamil Nadu

^c ICMR-National Institute of Epidemiology (ICMR-NIE), Chennai, Tamil Nadu

Corresponding author

Dr Meenakumari Ramaswamy

National Institute of Siddha (NIS),
Tambaram Sanatorium,
Chennai, Tamil Nadu 600047

Maid ID: mkumari474@yahoo.com